

# VAGUS NEET ACADEMY, TUMKUR

## MATHEMATICS KCET 2020 (VERSION CODE D2) KEY WITH SOLUTION

1. The sine of the angle between the straight line  $\frac{x-2}{3} = \frac{3-y}{-4} = \frac{z-4}{5}$  and the plane  $2x - 2y + z = 5$  is

(A)  $\frac{3}{50}$                       (B)  $\frac{4}{5\sqrt{2}}$                       (C)  $\frac{\sqrt{2}}{10}$                       (D)  $\frac{3}{\sqrt{50}}$

**Ans (C)**

Given line is  $\frac{x-2}{3} = \frac{y-3}{4} = \frac{z-4}{5}$

and plane is  $2x - 2y + z = 5$

w.k.t.  $\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| |\vec{n}|}$

$$\sin \theta = \frac{(3\hat{i} + 4\hat{j} + 5\hat{k}) \cdot (2\hat{i} - 2\hat{j} + \hat{k})}{\sqrt{9+16+25} \sqrt{4+4+1}}$$

$$= \frac{6-8+5}{\sqrt{50} \sqrt{9}} = \frac{3}{5\sqrt{2} \cdot 3} = \frac{1}{5\sqrt{2}} = \frac{1}{5\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{10}$$

2. If a line makes an angle of  $\frac{\pi}{3}$  with each of x and y-axis, then the acute angle made by z-axis is

(A)  $\frac{\pi}{6}$                       (B)  $\frac{\pi}{3}$                       (C)  $\frac{\pi}{2}$                       (D)  $\frac{\pi}{4}$

**Ans (D)**

Given  $\alpha = \beta = \frac{\pi}{3}$   $\gamma = ?$

$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$\left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2 + \cos^2 \gamma = 1$

$\frac{1}{4} + \frac{1}{4} + \cos^2 \gamma = 1$

$\cos^2 \gamma = \frac{1}{2} \Rightarrow \cos \gamma = \pm \frac{1}{\sqrt{2}}$

$\Rightarrow \gamma = \frac{\pi}{4}$  [ $\gamma$  is acute]

3. The distance of the point (1, 2, -4) from the line  $\frac{x-3}{2} = \frac{y-3}{3} = \frac{z+5}{6}$  is

(A)  $\frac{\sqrt{293}}{7}$                       (B)  $\frac{293}{49}$                       (C)  $\frac{\sqrt{293}}{49}$                       (D)  $\frac{293}{7}$

**Ans (A)**

Given point is (1, 2, -4) and the line is  $\frac{x-3}{2} = \frac{y-3}{3} = \frac{z+5}{6} = K$

$\frac{x-3}{2} = \frac{y-3}{3} = \frac{z+5}{6} = K$

$\therefore$  Dr's of the line are (2K + 3, 3K + 3, 6K - 5)


Dr's of AB  $(2K + 2, 3K + 1, 6K - 1)$

Since AB is perpendicular to the given line  $(2K + 2) \cdot 2 + (3K + 1) \cdot 3 + (6K - 1) \cdot 6 = 0$


$$4K + 4 + 9K + 3 + 36K - 6 = 0$$

$$49K = -1 \quad K = \frac{-1}{49}$$

$$\begin{aligned} \therefore \text{distance} &= \sqrt{\left(\frac{96}{49}\right)^2 + \left(\frac{46}{49}\right)^2 + \left(\frac{43}{49}\right)^2} \\ &= \sqrt{\frac{9216 + 2116 + 3025}{(49)^2}} = \frac{\sqrt{14357}}{\sqrt{49 \times 49}} = \frac{\sqrt{293}}{7} \end{aligned}$$



4. The feasible region of an LPP is shown in the figure. If  $Z = 11x + 7y$ , then the maximum value of  $Z$  occurs at



- (A)  $(3, 3)$                       (B)  $(5, 0)$                       (C)  $(3, 2)$                       (D)  $(0, 5)$

**Ans (C)**

The corner points are  $(0, 5), (0, 3), (3, 2)$

At  $(0, 5)$      $Z = 35$

At  $(0, 3)$      $Z = 21$

At  $(3, 2)$      $Z = 47$

5. Corner points of the feasible region determined by the system of linear constraints are  $(0, 3), (1, 1)$  and  $(3, 0)$ . Let  $z = px + qy$ , where  $p, q > 0$ . Condition on  $p$  and  $q$  so that the minimum of  $z$  occurs at  $(3, 0)$  and  $(1, 1)$  is

- (A)  $p = \frac{q}{2}$                       (B)  $p = 3q$                       (C)  $p = q$                       (D)  $p = 2q$

**Ans (A)**

Given corner points are  $(0, 3), (1, 1), (3, 0)$

$$z = px + qy$$

At  $(3, 0)$      $z = 3p$

At  $(1, 1)$      $z = p + q$

$$\Rightarrow 3p = p + q$$

$$2p = q \Rightarrow p = \frac{q}{2}$$

6. If A and B are two events such that  $P(A) = \frac{1}{3}, P(B) = \frac{1}{2}$  and  $P(A \cap B) = \frac{1}{6}$ , then  $P\left(\frac{A}{B}\right)$  is

- (A)  $\frac{1}{3}$                       (B)  $\frac{1}{2}$                       (C)  $\frac{1}{12}$                       (D)  $\frac{2}{3}$

**Ans (D)**

$$\text{Given } P(A) = \frac{1}{3} \quad P(B) = \frac{1}{2} \quad P(A \cap B) = \frac{1}{6}$$

$$P(A|B) = 1 - P(A|B)$$

$$= 1 - \frac{P(A \cap B)}{P(B)} = 1 - \frac{1}{3} = \frac{2}{3}$$

7. A die is thrown 10 times, the probability that an odd number will come up atleast one time is

- (A)  $\frac{1023}{1024}$                       (B)  $\frac{11}{1024}$                       (C)  $\frac{1013}{1024}$                       (D)  $\frac{1}{1024}$

**Ans (A)**

$$\text{Given } n = 10 \quad p = \frac{1}{2}, \quad q = \frac{1}{2}$$

$$\begin{aligned} \text{Required probability} &= 1 - P(X=0) \\ &= 1 - {}^{10}C_0 \left(\frac{1}{2}\right)^{10-0} \left(\frac{1}{2}\right)^0 \\ &= 1 - \frac{1}{2^{10}} = 1 - \frac{1}{1024} \\ &= \frac{1023}{1024} \end{aligned}$$

8. The probability of solving a problem by three persons A, B and C independently is  $\frac{1}{2}$ ,  $\frac{1}{4}$  and  $\frac{1}{3}$  respectively. Then the probability of the problem is solved by any two of them is

- (A)  $\frac{1}{4}$                       (C)  $\frac{1}{8}$                       —                      —

**Ans (A)**

(B)  $\frac{1}{24}$                       (D)  $\frac{1}{2}$

$$\begin{aligned} \text{Required probability} &= P(ABC) + P(ABC) + P(ABC) \\ &= \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{4} \cdot \frac{1}{3} \\ &= \frac{1}{12} + \frac{1}{12} + \frac{1}{12} = \frac{2+3+1}{24} = \frac{1}{4} \end{aligned}$$

9. Events  $E_1$  and  $E_2$  form a partition of the sample space S. A is any even such that  $P(E_1) = P(E_2) = \frac{1}{2}$ ,  $P(A|E_1) = \frac{1}{2}$  and  $P(A|E_2) = \frac{1}{3}$ , then  $P(E_1|A)$  is

- (A)  $\frac{2}{3}$                       (B) 1                      (C)  $\frac{1}{4}$                       (D)  $\frac{1}{2}$

**Ans (D)**

$$\begin{aligned} \text{Given } P(E_1) &= P(E_2) = \frac{1}{2} \\ P(E_1|A) &= \frac{1}{2} \quad P(A|E_2) = \frac{2}{3} \end{aligned}$$

$$\text{Using Baye's theorem } P(E_2|A) = \frac{P(E_2)P(A|E_2)}{P(E_1)P(A|E_1) + P(E_2)P(A|E_2)}$$

$$\frac{1}{2} = \frac{\binom{1}{2} \binom{2}{3}}{\binom{1}{2} x + \binom{1}{-} \binom{2}{\frac{1}{3}}} \Rightarrow \frac{1}{2} = \frac{3}{\frac{x}{2} + \frac{1}{3}}$$

$$\frac{x}{2} + \frac{1}{3} = \frac{2}{3} \Rightarrow \frac{x}{2} = \frac{1}{3} \Rightarrow x = \frac{2}{3}$$

$$P(E_1 | A) = \frac{P(E_1) P(A | E_1)}{P(E_1) P(A | E_1) + P(E_2) P(A | E_2)}$$

$$= \frac{\binom{1}{\frac{2}{3}} \binom{2}{3}}{\binom{1}{\frac{2}{3}} \binom{2}{3} + \binom{1}{-} \binom{2}{\frac{1}{3}}} = \frac{1}{2}$$

10. The value of  $\sin^2 51^\circ + \sin^2 39^\circ$  is

- (A) 0 (B)  $\sin 12^\circ$  (C)  $\cos 12^\circ$  (D) 1

**Ans (D)**

$$\sin^2 51^\circ + \sin^2 39^\circ = \cos^2 39^\circ + \sin^2 39^\circ = 1$$

11. If  $\tan A + \cot A = 2$ , then the value of  $\tan^4 A + \cot^4 A =$

- (A) 1 (B) 4 (C) 5 (D) 2

**Ans (D)**

$$\tan x + \cot x = 2$$

$$\tan^2 x + \cot^2 x + 2 \tan x \cot x = 2^2$$

$$\tan^2 x + \cot^2 x = 2$$

$$\tan^4 x + \cot^4 x + 2 \tan^2 x \cot^2 x = 4$$

$$\tan^4 x + \cot^4 x = 2$$

12. If  $A = \{1, 2, 3, 4, 5, 6\}$ , then the number of subsets of A which contain atleast two elements is

- (A) 63 (B) 57 (C) 58 (D) 64

**Ans (B)**

$$\text{Subsets of A are } 2^6 = 64$$

$$\text{Subsets of A which contain atleast two elements} = 64 - 7 = 57$$

13. If  $n(A) = 2$  and total number of possible relations from set A to set B is 1024, then  $n(B)$  is

- (A) 20 (B) 10 (C) 5 (D) 512

**Ans (C)**

$$n(A) = 2$$

$$2^m = 1024$$

$$(2^2)^n = 2^{10}$$

$$2^{2 \times 5} = 2^{10}$$

$$n(B) = 5$$

14. The value of

$${}^{16}C_9 + {}^{16}C_{10} - {}^{16}C_6 - {}^{16}C_7 \text{ is}$$

- (A) 1 (B)  ${}^{17}C_{10}$  (C)  ${}^{17}C_3$  (D) 0

**Ans (D)**

$${}^{16}C_9 + {}^{16}C_{10} - {}^{16}C_6 - {}^{16}C_7 = {}^{17}C_{10} - {}^{17}C_7 = {}^{17}C_{10} - {}^{17}C_{10} = 0$$

15. The number of terms in the expansion of  $(x + y + z)^{10}$  is  
 (A) 142 (B) 11 (C) 110 (D) 66

**Ans (D)**

$$\text{Number of terms in the expansion of } (x + y + z)^{10} = {}^{10+3-1}C_{10} = {}^{12}C_{10} = \frac{12!}{2!10!} = 66$$

16. If  $P(n) : 2^n < n!$   
 Then the smallest positive integer for which  $P(n)$  is true if  
 (A) 3 (B) 4 (C) 5 (D) 2

**Ans (B)**

$$P(n) : 2^n < n!$$

$$n = 4, 2^4 < 4!$$

$$n = 4$$

17. If  $z = x + iy$ , then the equation  $|z + 1| = |z - 1|$  represents  
 (A) a parabola (B) x-axis (C) y-axis (D) a circle

**Ans (C)**

$$|z + 1| = |z - 1|$$

$$|x + iy + 1| = |x + iy - 1|$$

$$\sqrt{(x+1)^2 + y^2} = \sqrt{(x-1)^2 + y^2}$$

$$(x+1)^2 + y^2 = (x-1)^2 + y^2$$

$$x^2 + 2x + 1 = x^2 + 1 - 2x$$

$$4x = 0$$

$$x = 0$$

y-axis

18. If the parabola  $x^2 = 4ay$  passes through the point  $(2, 1)$ , then the length of the latus rectum is  
 (A) 4 (B) 2 (C) 8 (D) 1

**Ans (A)**

$$x^2 = 4ay \quad \dots(1)$$

equation (1) passing through  $(2, 1)$

$$4 = 4a(1)$$

$$a = 1$$

$$\text{length of latus rectum} = 4a = 4(1) = 4$$

19. If the sum of  $n$  terms of an A.P. is given by  $S_n = n^2 + n$ , then the common difference of the A.P. is  
 (A) 1 (B) 2 (C) 6 (D) 4

**Ans (B)**

$$S_n = n^2 + n$$

$$S_1 = 1 + 1 = 2 = T_1$$

$$S_2 = 2^2 + 2 = 6 = T_1 + T_2$$

$$T_2 = S_2 - S_1 = 6 - 2 = 4$$

$$d = T_2 - T_1 = 4 - 2 = 2$$

20. The two lines  $lx + my = n$  and  $l'x + m'y = n'$  are perpendicular if  
 (A)  $lm' = ml$  (B)  $lm + l'm' = 0$  (C)  $lm' + ml' = 0$  (D)  $ll' + mm' = 0$

**Ans (D)**

The two lines  $lx + my = n$  and  $l'x + m'y = n'$  are perpendicular if  $ll' + mm' = 0$

21. The standard deviation of the data 6, 7, 8, 9, 10 is

- (A)  $\sqrt{10}$  (B) 2 (C) 10 (D)  $\sqrt{2}$

**Ans (D)**

$$\bar{x} = \frac{6 + 7 + 8 + 9 + 10}{5} = \frac{40}{5} = 8$$

$$\sigma = \sqrt{\frac{1}{5}[4 + 1 + 0 + 1 + 4]} \quad \square \quad \sigma = \sqrt{\frac{1}{n}(x_i - \bar{x})^2}$$

$$\sigma = \sqrt{\frac{10}{5}} = \sqrt{2}$$

22.  $\lim_{x \rightarrow 0} \left( \frac{\tan x}{\sqrt{2x+4}-2} \right)$  is equal to

- (A) 3 (B) 4 (C) 6 (D) 2

**Ans (D)**

$$\lim_{x \rightarrow 0} \left( \frac{\tan x}{\sqrt{2x+4}-2} \right) \text{ | L H' Rule}$$

$$= \lim_{x \rightarrow 0} \left( \frac{\sec^2 x}{\frac{1}{2 \times \sqrt{2x+4}} \times 2} \right)$$

$$= \lim_{x \rightarrow 0} (\sqrt{2x+4} \sec^2 x)$$

$$= \sqrt{2 \times 0 + 4} \times (\sec^2 0)$$

$$= 2 \times 1$$

$$= 2$$

23. The negation of the statement "For all real numbers x and y,  $x + y = y + x$ " is

- (A) for some real numbers x and y,  $x + y = y + x$   
(B) for some real numbers x and y,  $x + y \neq y + x$   
(C) for some real numbers x and y,  $x - y = y - x$   
(D) for all real numbers x and y,  $x + y \neq y + x$

**Ans (B)**

Negation : For some real numbers x and y,  $x + y \neq y + x$

24. Let  $f : [2, \infty) \rightarrow \mathbb{R}$  be the function defined by  $f(x) = x^2 - 4x + 5$ , then the range of f is

- (A)  $[1, \infty)$  (B)  $(1, \infty)$  (C)  $[5, \infty)$  (D)  $(-\infty, \infty)$

**Ans (A)**

$$\text{Let } f(x) = y$$

$$x^2 - 4x + 5 = y$$

$$x^2 - 2 \cdot x \cdot 2 + 4 + 1 = y$$

$$x^2 - 2 \cdot x \cdot 2 + 4 = y - 1$$

$$(x - 2)^2 = y - 1$$

$$x - 2 = \sqrt{y - 1}$$

$$x = \sqrt{y-1} + 2$$

$$\therefore y - 1 \geq 0$$

$$y \geq 1$$

Range is  $[1, \infty)$

25. If A, B, C are three mutually exclusive and exhaustive events of an experiment such that  $P(A) = 2P(B) = 3P(C)$ , then P(B) is equal to

$$(A) \frac{2}{11} \qquad \frac{P(A)}{2P(B)} = \frac{P(B)}{3} \qquad (B) \frac{11}{11}$$

$$(C) \frac{11}{11} \qquad (D) \frac{11}{11}$$

**Ans (B)**

$$P(A) = 2P(B) = 3P(C)$$

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$

$$1 = 2P(B) + P(B) + \frac{2}{3}P(B) - 0 - 0 - 0 + 0$$

$$1 = P(B) \left[ 3 + \frac{2}{3} \right]$$

$$1 = P(B) \left( \frac{11}{3} \right)$$

$$P(B) = \frac{3}{11}$$

26. If a relation R on the set  $\{1, 2, 3\}$  be defined by  $R = \{(1, 1)\}$ , then R is

- (A) Reflexive and transitive (B) Symmetric and transitive  
(C) Only symmetric (D) Reflexive and symmetric

**Ans (B)**

$$R = \{(1, 1)\} \text{ on a set } \{1, 2, 3\}$$

R is symmetric and Transitive

27. The value of  $\cos \left( \sin^{-1} \frac{\pi}{3} + \cos^{-1} \frac{\pi}{3} \right)$  is

- (A) 1 (B) -1 (C) Does not exist (D) 0

**Ans (D)**

$$\cos \left( \sin^{-1} \frac{\pi}{3} + \cos^{-1} \frac{\pi}{3} \right) = \cos \frac{\pi}{2} = 0$$

28. If  $A = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ , then  $A^4$  is equal to

- (A) 2A (B) I (C) 4A (D) A

**Ans (B)**

$$A^2 = AA = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = I$$

$$A^3 = A^2 \times A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = I$$

$$A^4 = A^3 A = I \times A = I$$

29. If  $A = \{a, b, c\}$ , then the number of binary operations on  $A$  is

- (A)  $3^6$  (B)  $3^3$  (C)  $3^9$  (D) 3

**Ans (C)**

$$A = \{a, b, c\}$$

The number of binary operations are  $n^{n^2} = 3^{3^2} = 3^9$

30. The domain of the function defined by  $f(x) = \cos^{-1} \sqrt{x-1}$  is

- (A)  $[0, 2]$  (B)  $[-1, 1]$  (C)  $[0, 1]$  (D)  $[1, 2]$

**Ans (D)**

$$f(x) = \cos^{-1} \sqrt{x-1}$$

$$-1 \leq \cos^{-1} x \leq 1$$

$$\text{and } -1 \leq \sqrt{x-1} \leq 1$$

$$0 \leq (x-1) \leq 1$$

$$1 \leq x \leq 2$$

31. If  $f(x) = \begin{vmatrix} x^3 - x & a + x & b + x \\ x - a & x^2 - x & c + x \\ x - b & x - c & 0 \end{vmatrix}$  then

- (A)  $f(2) = 0$  (B)  $f(0) = 0$  (C)  $f(-1) = 0$  (D)  $f(1) = 0$

**Ans (B)**

$$f(0) = \begin{vmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{vmatrix}$$

by [skew - symmetric matrix]

$$f(0) = 0$$

32. If  $A$  and  $B$  are square matrices of same order and  $B$  is a skew symmetric matrix, then  $ABA$  is

- (A) Null matrix (B) Diagonal matrix  
(C) Skew symmetric matrix (D) Symmetric matrix

**Ans (C)**

Given

$$B = -B'$$

$$\text{Now } (ABA)' = (BA)'(A)' \quad [\square (AB)' = BA']$$

$$= ABA$$

$$= -ABA$$



33. If  $\begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix} A = \begin{pmatrix} 1 & 0 \\ & \end{pmatrix}$ , then the matrix A is

- (A)  $\begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$  (B)  $\begin{pmatrix} -2 & 1 \\ 3 & -2 \end{pmatrix}$  (C)  $\begin{pmatrix} 2 & -1 \\ 3 & 2 \end{pmatrix}$  (D)  $\begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$

**Ans (A)**

We know that  $A^{-1}A = AA^{-1} = I$  or  $BA = AB = I$

$$\text{Let } B = \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix} \Rightarrow |B| = 4 - 3 = 1$$

$$\text{adj}B = \begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$$

$$B^{-1} = \begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$$

$$\therefore A = B^{-1} = \begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$$

34. If  $f(x) = \begin{cases} \frac{1 - \cos Kx}{x \sin x} & , \text{ if } x \neq 0 \\ \frac{1}{2} & , \text{ if } x = 0 \end{cases}$  is continuous at  $x = 0$ , then the value of K is

- (A) 0 (B)  $\pm 2$  (C)  $\pm 1$  (D)  $\pm \frac{1}{2}$

**Ans (C)**

Given

$$\lim_{x \rightarrow 0} f(x) = f(0)$$

$$\lim_{x \rightarrow 0} \left( \frac{1 - \cos Kx}{x \sin x} \right) = \frac{1}{2}$$

By L'H rule

$$\lim_{x \rightarrow 0} \left( \frac{\sin Kx \cdot K}{x \cos x + \sin x} \right) = \frac{1}{2}$$

By LH rule

$$\lim_{x \rightarrow 0} \left( \frac{\cos Kx \cdot K^2}{-x \sin x + \cos x + \cos x} \right) = \frac{1}{2}$$

$$\frac{1 \cdot K^2}{0 + 1 + 1} = \frac{1}{2} \Rightarrow K^2 = 1$$

$$K = \pm 1$$

35. If  $a_1, a_2, a_3, \dots, a_9$  are in A.P. then the value of  $\begin{vmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{vmatrix}$  is

- (A)  $a_1 + a_9$  (B)  $\log_e(\log_e e)$  (C) 1 (D)  $\frac{9}{2} (a_1 + a_9)$

**Ans (B)**

Let a be first term and d be common difference

$$GE = \begin{vmatrix} a & a+d & a+2d \\ a+3d & a+4d & a+5d \\ a+6d & a+7d & a+8d \end{vmatrix}$$

by applying  $C_2 \rightarrow C_2 - C_1$

$$C_3 \rightarrow C_3 - C_2$$

$$= \begin{vmatrix} a & d & d \\ a+3d & d & d \\ a+6d & d & d \end{vmatrix}$$

$$= 0$$

$$= \log(\log_e e) [\square \log 1 = 0]$$

36. If A is a square matrix of order 3 and  $|A| = 5$ , then  $|A \text{ adj } A|$  is

(A) 125

(B) 25

(C) 625

(D) 5

**Ans (A)**

$$|A \text{ adj } A| = |A| |\text{adj } A|$$

$$= |A| |A|^{3-1}$$

$$= 5 \cdot 5^2$$

$$= 125$$

37. If  $f(x) = \sin^{-1} \left( \frac{2x}{\sqrt{1+x^2}} \right)$ , then  $f(\sqrt{3})$  is

(A)  $\frac{1}{2}$

(B)  $\left(\frac{1}{\sqrt{3}}\right)$

(C)  $-\frac{1}{\sqrt{3}}$

(D)  $-\frac{1}{2}$

**Ans (A)**

$$\text{Put } x = \tan \theta \Rightarrow \theta = \tan^{-1} x$$

$$f(x) = \sin^{-1} \left( \frac{2 \tan \theta}{\sqrt{1 + \tan^2 \theta}} \right)$$

$$= \sin^{-1} (\sin 2\theta)$$

$$= 2\theta$$

$$f(x) = 2 \tan^{-1} x$$

$$f(x) = \frac{2}{1+x^2}$$

$$f(\sqrt{3}) = \frac{2}{1+3} = \frac{1}{2}$$

38. The right hand and left hand limit of the function  $f(x) = \begin{cases} e^{1/x} - 1 & , \text{ if } x \neq 0 \\ 0 & , \text{ if } x = 0 \end{cases}$  are respectively

(A) 1 and -1

(B) -1 and -1

(C) -1 and 1

(D) 1 and 1

**Ans (D)**

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left( \frac{e^{1/x} - 1}{\frac{1}{e^x + 1}} \right)$$

By dividing both numerator and denominator by  $e^{\frac{1}{x}}$

$$\begin{aligned}
 &= \lim_{x \rightarrow 0} \left( \frac{1 - e^{-\frac{1}{x}}}{1 + e^{-\frac{1}{x}}} \right) \\
 &= \lim_{x \rightarrow 0} \left( \frac{1 - 0}{1 + 0} \right) \left[ \begin{array}{l} \square e^{\infty} = \infty \\ -\infty \quad 1 \\ e^{-\frac{1}{\infty}} = 0 \end{array} \right] \\
 &= 1
 \end{aligned}$$

$\therefore \text{LHL} = \text{RHL} = 1$

39. If  $2^x + 2^y = 2^{x+y}$ , then  $\frac{dy}{dx}$  is

(A)  $-2^{y-x}$

(B)  $2^{x-y}$

(C)  $\left( \frac{2^y - 1}{2^x - 1} \right)$

(D)  $2^{y-x}$

**Ans (A)**

We know that,

If  $a^x + a^y = a^{x+y}$  then  $\frac{dy}{dx} = -a^{y-x}$

$\therefore \frac{dy}{dx} = -2^{y-x}$

40. If the curves  $2x = y^2$  and  $2xy = k$  intersect perpendicularly, then the value of  $k^2$  is

(A)  $2\sqrt{2}$

(B) 2

(C) 8

(D) 4

**Ans (C)**

$2x = y^2 \dots(1)$

$2xy = k \dots(2)$

Solve (1) and (2)

$(2) \Rightarrow y^3 = k$

$y = k^{\frac{1}{3}}$

$(1) \Rightarrow x = \frac{y^2}{2} = \frac{k^{\frac{2}{3}}}{2}$

$(x, y) = \left( \frac{k^{\frac{2}{3}}}{2}, k^{\frac{1}{3}} \right)$

Differentiate (1) w.r.t x

$y' = \frac{1}{y}$

$m_1 = \frac{1}{\frac{1}{k^{\frac{1}{3}}}}$

Differentiate (2) w.r.t x

$y' = -\frac{y}{x}$

$m_2 = \frac{\frac{1}{2k^{\frac{2}{3}}}}{\frac{1}{k^{\frac{1}{3}}}} = \frac{-2}{k^{\frac{1}{3}}}$

Given

$$m_1 m_2 = -1$$

$$\frac{1}{-2} \times \frac{1}{1} = -1$$

$$k^3 = k^3$$

$$\frac{2}{k^3} = 2$$

$$k^2 = 8$$

41. If  $(xe)^y = e^x$ , then  $\frac{dy}{dx}$  is

(A)  $\frac{1}{(1+\log x)^2}$

(B)  $\frac{\log x}{(1+\log x)}$

(C)  $\frac{e^x}{x(y-1)}$

(D)  $\frac{\log x}{(1+\log x)^2}$

**Ans (D)**

$$(xe)^y = e^x$$

$$\Rightarrow y(\log x + 1) = x$$

$$\Rightarrow y = \frac{x}{\log x + 1}$$

$$\therefore \frac{dy}{dx} = \frac{\log x}{(\log x + 1)^2}$$

42. If  $y = 2x^{n+1} + \frac{3}{x^n}$ , then  $x^2 \frac{d^2y}{dx^2}$  is

(A)  $n(n+1)y$

(B)  $x \frac{dy}{dx} + y$

(C)  $y$

(D)  $6n(n+1)y$

**Ans (A)**

$$y = 2x^{n+1} + 3x^{-n}$$

$$\Rightarrow \frac{dy}{dx} = 2(n+1)x^n - 3nx^{-n-1}$$

$$\Rightarrow \frac{d^2y}{dx^2} = 2n(n+1)x^{n-1} + 3n(n+1)x^{-n-2}$$

$$\Rightarrow x^2 \frac{d^2y}{dx^2} = n(n+1) \left[ 2x^{n+1} + \frac{3}{x^n} \right]$$

$$\Rightarrow x^2 \frac{d^2y}{dx^2} = n(n+1)y$$

43. The value of  $\int \frac{1+x^4}{1+x^6} dx$  is

(A)  $\tan^{-1} x + \frac{1}{3} \tan^{-1} x^3 + C$

(B)  $\tan^{-1} x - \frac{1}{3} \tan^{-1} x^3 + C$

(C)  $\tan^{-1} x + \frac{1}{3} \tan^{-1} x^2 + C$

(D)  $\tan^{-1} x + \tan^{-1} x^3 + C$

**Ans (A)**

$$\int \frac{1+x^4}{1+x^6} dx = \int \frac{1+x^4-x^2+x^2}{(1+x^2)(1-x^2+x^4)} dx$$

$$= \int \frac{1-x^2+x^4}{(1+x^2)(1-x^2+x^4)} + \frac{x^2}{(1+x^2)(1-x^2+x^2)} dx$$

$$\begin{aligned}
 &= \int \left( \frac{1}{1+x^2} + \frac{x^2}{1+x^6} \right) dx \\
 &= \int \left( \frac{1}{1+x^2} + \frac{1}{3} \frac{3x^2}{1+(x^3)^2} \right) dx \\
 &= \tan^{-1} x + \frac{1}{3} \tan^{-1}(x^3) + C
 \end{aligned}$$

44. The maximum value of  $\frac{\log_e x}{x}$ , if  $x > 0$  is

- (A) 1                                      (B)  $\frac{1}{e}$                                       (C)  $-\frac{1}{e}$                                       (D) e

**Ans (B)**

$$\begin{aligned}
 y &= \frac{\log x}{x} \\
 \Rightarrow \frac{dy}{dx} &= \frac{1 - \log x}{x^2} \\
 \frac{dy}{dx} &= 0 \\
 \Rightarrow 1 - \log x &= 0 \\
 \Rightarrow x &= e \\
 \therefore y_{\max} &= \frac{1}{e}
 \end{aligned}$$

45. If the side of a cube is increased by 5%, then the surface area of a cube is increased by

- (A) 60%                                      (B) 6%                                      (C) 20%                                      (D) 10%

**Ans (D)**

$$\begin{aligned}
 A &= 6x^2 & \frac{dx}{dt} &= \frac{5x}{100} \\
 \frac{dA}{dt} &= 12x \frac{dx}{dt} = 12x \times \frac{5x}{100} = \frac{60x^2}{100} \\
 \Rightarrow &= \frac{10}{100} \times 6x^2 = \frac{10}{100} A \\
 \therefore &10\%
 \end{aligned}$$

46. The value of  $\int_{-\frac{1}{2}}^{\frac{1}{2}} \cos^{-1} x \, dx$  is

- (A)  $\frac{\pi}{2}$                                       (B) 1                                      (C)  $\frac{\pi^2}{2}$                                       (D)  $\pi$

**Ans (A)**

$$\begin{aligned}
 \int_{-\frac{1}{2}}^{\frac{1}{2}} \cos^{-1} x \, dx &= x \cos^{-1} x \Big|_{-\frac{1}{2}}^{\frac{1}{2}} + \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{x}{\sqrt{1-x^2}} dx \\
 &= \frac{1}{2} \cos^{-1} \left( \frac{1}{2} \right) - \left( -\frac{1}{2} \right) \cos^{-1} \left( -\frac{1}{2} \right) - \frac{1}{2} \cdot 2 \sqrt{1-x^2} \Big|_{-\frac{1}{2}}^{\frac{1}{2}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \cdot \frac{\pi}{3} + \frac{1}{2} \cdot \frac{2\pi}{3} - 0 \\
 &= \frac{\pi}{6} + \frac{\pi}{3} = \frac{\pi}{2}
 \end{aligned}$$

47. If  $\int \frac{3x+1}{(x-1)(x-2)(x-3)} dx = A \log|x-1| + B \log|x-2| + C \log|x-3| + C$ , then the values of A, B and C are respectively,

- (A) 2, -7, -5                      (B) 5, -7, 5                      (C) 2, -7, 5                      (D) 5, -7, -5

**Ans (C)**

$$\frac{3x+1}{(x-1)(x-2)(x-3)} = \frac{A}{x-1} + \frac{B}{x-2} + \frac{C}{x-3}$$

$$\Rightarrow 3x+1 = A(x-2)(x-3) + B(x-1)(x-3) + C(x-1)(x-2)$$

$$x=1 \Rightarrow A=2, x=2 \Rightarrow B=-7, x=3 \Rightarrow C=5$$

48. The value of  $\int e^{\sin x} \sin 2x dx$  is

- (A)  $2 e^{\sin x} (\sin x + 1) + C$                       (B)  $2 e^{\sin x} (\cos x + 1) + C$   
 (C)  $2 e^{\sin x} (\cos x - 1) + C$                       (D)  $2 e^{\sin x} (\sin x - 1) + C$

**Ans (D)**

$$\int e^{\sin x} \sin 2x dx = 2 \int e^{\sin x} \sin x \cos x dx = 2 \int t e^t dt$$

$$= 2 [te^t - e^t] + c$$

$$\sin x = t$$

$$= 2 [\sin x - 1] e^{\sin x} + c$$

$$\Rightarrow \cos x dx = dt$$

49. The area of the region bounded by the curve  $y^2 = 8x$  and the line  $y = 2x$  is

- (A)  $\frac{4}{3}$  sq. units                      (B)  $\frac{3}{4}$  sq. units                      (C)  $\frac{8}{3}$  sq. units                      (D)  $\frac{16}{3}$  sq. units

**Ans (A)**

$$y^2 = 8x \text{ and } y = 2x$$

$$\Rightarrow 4x^2 = 8x$$

$$\Rightarrow x^2 - 2x = 0$$

$$\Rightarrow x = 0, 2$$

$$RA = \int_0^2 |8\sqrt{x} - 2x| dx$$

$$= 2 \sqrt{2} \cdot \left[ \frac{x^{3/2}}{3/2} - x^2 \right]_0^2 = \frac{4\sqrt{2}}{3} (2^{3/2} - 2^2 - 0)$$

$$= \frac{4\sqrt{2}}{3} \cdot 2\sqrt{2} - 4 = \frac{16}{3} - 4$$

$$= \frac{4}{3} \text{ sq. units}$$

50. The value of  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos x}{1+e^x} dx$  is

- (A) 0 (B) 1 (C) -2 (D) 2

**Ans (B)**

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos x}{1+e^x} dx \quad \dots(1)$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos\left(\frac{\pi}{2} - \frac{\pi}{2} - x\right)}{1+e^{-x}} dx$$

$$I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{-\cos x}{1+e^{-x}} dx = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{e^x \cos x}{-1+e^x} dx \quad \dots(2)$$

$$(1) + (2) \Rightarrow 2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{(e^x + 1) \cos x}{e^x + 1} dx$$

$$2I = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos x dx = \sin x \Big|_{-\frac{\pi}{2}}^{\frac{\pi}{2}}$$

$$2I = \sin \frac{\pi}{2} - \sin\left(-\frac{\pi}{2}\right)$$

$$2I = 1 - (-1) = 2$$

$$\therefore I = 1$$

51. The value of  $\int_0^1 \frac{\log(1+x)}{1+x^2} dx$  is

- (A)  $\frac{\pi}{4} \log 2$  (B)  $\frac{1}{2}$  (C)  $\frac{\pi}{8} \log 2$  (D)  $\frac{\pi}{2} \log 2$

**Ans (C)**

$$\int_0^1 \frac{\log(1+x)}{1+x^2} dx \quad \left| \begin{array}{l} x = \tan \theta \\ dx = \sec^2 \theta d\theta \end{array} \right.$$

$$= \int_0^{\frac{\pi}{4}} \log(1+\tan \theta) d\theta$$

$$= \frac{\pi}{8} \log 2$$

52. The general solution of the differential equation  $x^2 dy - 2xy dx = x^4 \cos x dx$  is

- (A)  $y = x^2 \sin x + c$  (B)  $y = \sin x + cx^2$   
 (C)  $y = \cos x + cx^2$  (D)  $y = x^2 \sin x + cx^2$

**Ans (D)**

$$x^2 dy - 2xy dx = x^4 \cos x dx$$

$$\frac{dy}{dx} = \frac{x^4 \cos x + 2xy}{x^2}$$

$$\frac{dy}{dx} - \frac{2}{x}y = x^2 \cos x$$

$$\text{IF} = e^{\int -\frac{2}{x} dx} = e^{-2 \log x} = \frac{1}{x^2}$$

$$\text{General solution is } y \left( \frac{1}{x^2} \right) = \int \frac{1}{x^2} (x^2 \cos x) dx + c$$

$$\frac{y}{x^2} = \sin x + c$$

53. The area of the region bounded by the line  $y = 2x + 1$ , x-axis and the ordinates  $x = -1$  and  $x = 1$  is


(A) 2

(B)  $\frac{5}{2}$

(C) 5

(D)  $\frac{9}{4}$

**Ans (B)**



$$\begin{aligned} \text{Area bounded by } y = 2x + 1 \text{ with x-axis} &= \frac{1}{2} \begin{pmatrix} 1 \\ -1/2 \end{pmatrix} (1) + \frac{1}{2} \begin{pmatrix} 3 \\ 1/2 \end{pmatrix} (3) \\ &= \frac{1}{4} + \frac{9}{4} \\ &= \frac{10}{4} \\ &= \frac{5}{2} \text{ sq. units} \end{aligned}$$

54. The order of the differential equation obtained by eliminating arbitrary constants in the family of curves

$$c_1 y = (c_2 + c_3) e^{x+c_4} \text{ is}$$

(A) 2

(B) 3

(C) 4

(D) 1

**Ans (D)**

$$c_1 y = (c_2 + c_3) e^{x+c_4}$$

$$y = \left( \frac{c_2 + c_3}{c_1} e^{c_4} \right) e^x$$

$$y = Ae^x$$

Order = number of arbitrary constants

$$= 1$$



55. If  $\vec{a}$  and  $\vec{b}$  are unit vectors and  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ , then  $\sin \frac{\theta}{2}$  is

- (A)  $\frac{|\vec{a} + \vec{b}|}{2}$                       (B)  $\frac{|\vec{a} - \vec{b}|}{2}$                       (C)  $|\vec{a} - \vec{b}|$                       (D)  $|\vec{a} + \vec{b}|$

**Ans (B)**

$$|\vec{a} - \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}||\vec{b}|\cos\theta$$

(  $|\vec{a}| = |\vec{b}| = 1$  )

$$|\vec{a} - \vec{b}|^2 = 2\left(1 - \cos\theta\right)$$

$$\sin \frac{\theta}{2} = \frac{|\vec{a} - \vec{b}|}{2}$$

56. The curve passing through the point (1, 2) given that the slope of the tangent at any point (x, y) is  $\frac{2x}{y}$

represents

- (A) Parabola                      (B) Ellipse                      (C) Hyperbola                      (D) Circle

**Ans (C)**

$$\text{Slope} = \frac{dy}{dx} = \frac{2x}{y}$$

$$\Rightarrow y \, dy = 2x \, dx$$

$$\Rightarrow \int y \, dy = \int 2x \, dx + A$$

$$\Rightarrow \frac{y^2}{2} = x^2 + A \Rightarrow \frac{y^2}{2} - x^2 = A$$

$\Rightarrow$  curve is hyperbola

57. The two vectors  $\hat{i} + \hat{j} + \hat{k}$  and  $\hat{i} + 3\hat{j} + 5\hat{k}$  represent the two sides AB and AC respectively of a  $\Delta ABC$ . The length of the median through A is

- (A) 14                      (B) 7                      (C)  $\sqrt{14}$                       (D)  $\frac{\sqrt{14}}{2}$

**Ans (C)**

$$\vec{AB} = (1, 1, 1) \quad \vec{AC} = (1, 3, 5)$$

$$\vec{BC} = (0, 2, 4)$$


$$\vec{BD} = (0, 1, 2)$$

$$\vec{AD} = \vec{AB} + \vec{BD}$$

$$= (1, 2, 3)$$

$$|\vec{AD}| = \sqrt{1+4+9}$$

$$= \sqrt{14}$$



58. The point (1, -3, 4) lies in the octant

- (A) Third                      (B) Fourth                      (C) Eighth                      (D) Second

**Ans (B)**

$$(1, -3, 4)$$

Fourth octant

59. If the vectors  $2\hat{i} - 3\hat{j} + 4\hat{k}$ ,  $2\hat{i} + \hat{j} - \hat{k}$  and  $\lambda\hat{i} - \hat{j} + 2\hat{k}$  are coplanar, then the value of  $\lambda$  is  
 (A) -5 (B) -6 (C) 5 (D) 6

**Ans (D)**

Vectors are coplanar

$$\Rightarrow \begin{vmatrix} 2 & -3 & 4 \\ 2 & 1 & -1 \\ \lambda & -1 & 2 \end{vmatrix} = 0$$

$$\Rightarrow 2(1) + 3(4 + \lambda) + 4(-2 - \lambda) = 0$$

$$\Rightarrow 2 + 12 + 3\lambda - 8 - 4\lambda = 0$$

$$\Rightarrow \lambda = 6$$

60. If  $|\mathbf{a} \times \mathbf{b}| + \mathbf{a} \cdot \mathbf{b} = 144$  and  $|\mathbf{a}| = 6$ , then  $|\mathbf{b}|$  is equal to

(A) 3

(B) 2

(C) 4

(D) 6

**Ans (B)**

$$|\mathbf{a} \times \mathbf{b}| + \mathbf{a} \cdot \mathbf{b} = 144$$

$$\Rightarrow |\mathbf{a}| |\mathbf{b}| \sin \theta + |\mathbf{a}| |\mathbf{b}| \cos \theta = 144$$

$$\Rightarrow |\mathbf{b}| = \frac{144}{36} = 4$$

$$\Rightarrow |\mathbf{b}| = 2$$

\* \* \*